

Anti-PBEF / Visfatin / NAMPT antibody



Catalog Number: 100491

Product name

Anti-PBEF / Visfatin / NAMPT antibody

Immunogen

[Human PBEF / Visfatin / NAMPT \(His & GST Tag\) recombinant protein](#)

Specificity

Human NAMPT / PBEF

Antibody description

Rabbit polyclonal to PBEF / Visfatin / NAMPT

Preparation

Produced in rabbits immunized with purified, recombinant Human NAMPT / PBEF (rh NAMPT / PBEF; P43490; Met1-His491). NAMPT / PBEF specific IgG was purified by Human NAMPT / PBEF affinity chromatography.

Formulation

0.2 μ m filtered solution in PBS with 5% trehalose

Storage

This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C.

Preservative-Free.

Sodium azide is recommended to avoid contamination (final concentration 0.05%-0.1%). It is toxic to cells and should be disposed of properly. Avoid repeated freeze-thaw cycles.

Clonality

Polyclonal

Ig Type

Rabbit IgG

Applications

ELISA, WB, IHC-P, IP

Dilutions

WB: 5-10 μ g/ml


ELISA: 0.1-0.2 μ g/mL

This antibody can be used at 0.1-0.2 μ g/mL with the appropriate secondary reagents to detect Human NAMPT / PBEF. The detection limit for Human NAMPT / PBEF is approximately 0.00975 ng/well.


IHC-P: 0.1-2 μ g/mL

IP: 1-4 μ g/mg of lysate

Validations




PBEF / Visfatin / NAMPT Antibody, Rabbit PAb, Antigen Affinity Purified



PBEF / Visfatin / NAMPT Antibody, Rabbit PAb, Antigen Affinity Purified, Immunohistochemistry

Catalog Number: 100491


Immunochemical staining of human NAMPT in human kidney with rabbit polyclonal antibody (1 $\mu\text{g/mL}$, formalin-fixed paraffin embedded



sections).

PBEF / Visfatin / NAMPT Antibody, Rabbit PAb, Antigen Affinity Purified, Immunohistochemistry

Immunochemical staining of human NAMPT in human liver with rabbit polyclonal antibody (1 $\mu\text{g/mL}$, formalin-fixed paraffin embedded



sections).

PBEF / Visfatin / NAMPT Antibody, Rabbit PAb, Antigen Affinity Purified, Western blot

Lanes	
Items	A
Sample (whole cell lysate)	Jurkat
Sample Volume ($\mu\text{g/lane}$)	30
Gel	13% SDS-PAGE reducing gel
Recommended Concentration	5-10 $\mu\text{g/ml}$
Secondary Antibody	Dylight 800 labeled Antibody To Rabbit IgG (H+L), at 1:5000 dilution.