

Anti-FOS / c-Fos antibody



Catalog Number: 100670

Product name

Anti-FOS / c-Fos antibody

IHC-P: 0.1-2 µg/mL

ICC/IF: 0.5-1.5 µg/mL

Specificity

Human FOS / c-Fos

Antibody description

Rabbit polyclonal to FOS / c-Fos

Preparation

Produced in rabbits immunized with a synthetic peptide corresponding to the center region of the Human FOS / c-Fos, and purified by antigen affinity chromatography.

Formulation

0.2 µm filtered solution in PBS

Storage

This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C.

Preservative-Free.

Sodium azide is recommended to avoid contamination (final concentration 0.05%-0.1%). It is toxic to cells and should be disposed of properly. Avoid repeated freeze-thaw cycles.

Clonality

Polyclonal

Ig Type


Rabbit IgG

Applications

IHC-P, IF, ICC/IF


Dilutions

Validations



FOS / c-Fos Antibody, Rabbit PAb, Antigen Affinity Purified, Immunohistochemistry

Immunochemical staining of human FOS in human brain with rabbit polyclonal antibody (0.5 µg/mL, formalin-fixed paraffin embedded sections).




FOS / c-Fos Antibody, Rabbit PAb, Antigen Affinity Purified, Immunohistochemistry

Immunochemical staining of human FOS in human breast carcinoma with rabbit polyclonal antibody (0.5 µg/mL, formalin-fixed paraffin embedded

Anti-FOS / c-Fos antibody




Catalog Number: 100670



sections).

FOS / c-Fos Antibody, Rabbit PAb, Antigen Affinity Purified, Immunohistochemistry

Immunochemical staining of human FOS in human skin with rabbit polyclonal antibody (0.5 µg/mL, formalin-fixed paraffin embedded sections).



FOS / c-Fos Antibody, Rabbit PAb, Antigen Affinity Purified, Immunofluorescence

Immunofluorescence staining of FOS in HeLa cells. Cells were fixed with 4% PFA, permeabilized with 0.3% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-human FOS polyclonal antibody (1 µg/ml) at 4°C overnight. Then cells were stai