

# Anti-BLNK/Ly-57/SLP-65 antibody



Catalog Number: 103370

## Product name

Anti-BLNK/Ly-57/SLP-65 antibody

## Immunogen

[Human BLNK/Ly-57/SLP-65 \(His Tag\) recombinant protein](#)

## Specificity

Human BLNK / Ly-57 / SLP-65

## Antibody description

Rabbit polyclonal to BLNK/Ly-57/SLP-65

## Preparation

Produced in rabbits immunized with purified, recombinant Human BLNK / Ly-57 / SLP-65 (rh BLNK / Ly-57 / SLP-65; P08195-1; Met 1-Ser 456). BLNK / Ly-57 / SLP-65 specific IgG was purified by Human BLNK / Ly-57 / SLP-65 affinity chromatography.

## Formulation

0.2 µm filtered solution in PBS with 5% trehalose

## Storage

This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C.

Preservative-Free.

Sodium azide is recommended to avoid contamination (final concentration 0.05%-0.1%). It is toxic to cells and should be disposed of properly. Avoid repeated freeze-thaw cycles.

## Clonality

Polyclonal

## Ig Type

Rabbit IgG

## Applications

ELISA, WB, IHC-P

## Dilutions


WB: 10-20 µg/mL

ELISA: 0.1-0.2 µg/mL

This antibody can be used at 0.1-0.2 µg/mL with the appropriate secondary reagents to detect Human BLNK. The detection limit for Human BLNK is approximately 0.039 ng/well.


IHC-P: 0.1-2 µg/mL

## Validations



BLNK / Ly-57 / SLP-65 Antibody, Rabbit PAb, Antigen Affinity Purified, Immunohistochemistry

Immunochemical staining of human BLNK in human brain with rabbit polyclonal antibody (1 µg/mL, formalin-fixed paraffin embedded



sections).


Anti-BLNK/Ly-57/SLP-65 antibody



Catalog Number: 103370

BLNK / Ly-57 / SLP-65 Antibody, Rabbit PAb,  
Antigen Affinity Purified, Immunohistochemistry

Immunochemical staining of human BLNK in  
human spleen with rabbit polyclonal antibody (1  
µg/mL, formalin-fixed paraffin embedded




sections).

BLNK / Ly-57 / SLP-65 Antibody, Rabbit PAb,

Antigen Affinity Purified, Immunohistochemistry

Immunochemical staining of human BLNK in  
human tonsil with rabbit polyclonal antibody (1  
µg/mL, formalin-fixed paraffin embedded



Lanes	
Items	A
Sample {whole cell lysate }	Raji
Sample Volume (µg/lane)	30
Gel	12%SDS-PAGE reducing gel
Recommended Concentration	10-20 µg/ml
Secondary Antibody	Dylight 800-labeled Antibody To Rabbit IgG (H+L), at 1:5000 dilution.
Developed using Odyssey imaging system.	
Explanation	Predicted band size : 50 kDa Observed band size : 70 kDa

sections).

BLNK / Ly-57 / SLP-65 Antibody, Rabbit PAb,  
Antigen Affinity Purified, Western blot