

Anti-TAGLN antibody



Catalog Number: 142802

Product name

Anti-TAGLN antibody

WB: 5-20 µg/ml

IHC-P: 0.1-2µg/mL

Specificity

Human TAGLN

ICC/IF: 0.5-3µg/mL

IP: 4-8 uL/mg of lysate


Antibody description

Rabbit Polyclonal to Human TAGLN

Validations

Preparation

Produced in rabbits immunized with a synthetic peptide corresponding to the C-terminus of the Human TAGLN, and purified by antigen affinity chromatography.



Formulation

0.2 µm filtered solution in PBS


Human TAGLN Immunofluorescence(IF) 15569

Storage

This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C.

Preservative-Free.

Sodium azide is recommended to avoid contamination (final concentration 0.05%-0.1%). It is toxic to cells and should be disposed of properly. Avoid repeated freeze-thaw cycles.



Item	Lanes	A
Sample (whole cell lysate)		Jurkat
Sample quantity		0.5 mg
IP antibody quantity		4 µg
Protein G agarose		15 µl of 50% Protein G agarose
Gel		13% SDS-PAGE reducing gel
Primary antibody		KLH-SMCC-sinoA2084/KLH-SMCC-sinoA2141 antibody at 10 µg/ml
Secondary antibody		DyLight 800-labeled antibody to rabbit IgG (H+L), at 1:5000 dilution

Human TAGLN Immunoprecipitation(IP) 15570

Clonality


Polyclonal

Ig Type

Rabbit IgG

Applications

WB, IHC-P, ICC/IF, IF, IP




Human TAGLN Immunohistochemistry(IHC)

Dilutions


Anti-TAGLN antibody

Catalog Number: 142802



15568

Human TAGLN Immunohistochemistry(IHC)



Items	Lanes	A
Sample (whole cell lysate)		Hela
Sample Volume (µg/lane)		30
Gel		13% SDS-PAGE reducing gel
Recommended Concentration		5-20 µg/ml
Secondary Antibody		Dylight 800-labeled Antibody to Rabbit IgG (H+L), at 1:5000 dilution.
		Developed using Odyssey imaging system.
Explanation		Predicted band size : 23 kDa Observed band size : 23 kDa

15567

Human TAGLN Western blot (WB) 15566