

Product name

Anti-Selp antibody

Specificity

Human, Mouse, Rat, Dog, Pig, Cow, Horse

Antibody description

Rabbit polyclonal antibody to Selp

Preparation

This antigen of this antibody was klh conjugated synthetic peptide derived from mouse p-selectin 701-768/768

Formulation

Liquid, 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

Storage

Store at -20°C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4°C.

Clonality

Polyclonal

Ig Type

IgG

Applications

WB, IHC-P, FC

Dilutions

IHC-P:1:400-800

FC:1µg/Test

Validations




Fig1: Blank control: HUVEC cells(blue).; Primary Antibody:Rabbit Anti-CD62p antibody , Dilution: 1µg in 100 µL 1X PBS containing 0.5% BSA;; Isotype Control Antibody: Rabbit IgG(orange) ,used under the same conditions);; Secondary Antibody: Goat anti-rabbit IgG-PE(white blue), Dilution: 1:200 in 1 X PBS containing 0.5% BSA.; Protocol; The cells were fixed with 2% paraformaldehyde (10 min) .Primary antibody (1µg /1x10⁶ cells) were incubated for 30 min on the ice, followed by 1 X PBS containing 0.5% BSA + 1 0% goat serum (15 min) to block non-specific protein-protein interactions. Then the Goat Anti-rabbit IgG/PE antibody was added into the blocking buffer mentioned above to react with the primary antibody at 1/200 dilution for 30 min on ice. Acquisition of 20,000 events was performed.




Fig2: Tissue/cell: rat pancreas tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min;

Anti-Selp antibody



Catalog Number: 175319

Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;; Incubation: Anti-P-selectin Polyclonal Antibody, Unconjugated 1:200,

overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining