

Anti-NDUFB8 antibody

Catalog Number: 176582

Product name

Anti-NDUFB8 antibody

Specificity

Human, Mouse, Rat

Antibody description

Rabbit monoclonal antibody to NDUFB8

Preparation

This antigen of this antibody was recombinant protein within n-terminal human ndufb8 .

Formulation

Liquid, 1*TBS (pH7.4), 0.05% BSA, 40% Glycerol.
Preservative: 0.05% Sodium Azide.

Storage

Store at +4°C after thawing. Aliquot store at -20°C. Avoid repeated freeze / thaw cycles.

Clonality

Monoclonal

Ig Type

IgG

Applications

WB, IP, IHC-P, FC, ICC

Dilutions

WB: 1:500-1:2,000

IP: 1:10-1:50

IHC-P: 1:50-1:200

FC: 1:50-1:100

ICC: 1:50

Validations




Fig1:: Western blot analysis of NDUFB8 on different lysates. Proteins were transferred to a PVDF membrane and blocked with 5% BSA in PBS for 1 hour at room temperature. The primary antibody (1/500) was used in 5% BSA at room temperature for 2 hours. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1:40,000 dilution was used for 1 hour at room temperature.; Positive control:; Lane 1: 293 cell lysate; Lane 2: A549 cell lysate; Lane 3: Mouse heart tissue lysate; Lane 4: Rat spleen tissue lysate




Fig2:: Immunohistochemical analysis of paraffin-embedded rat heart tissue using anti-NDUFB8 antibody. The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 8.0-8.4) for 20 minutes. The tissues were blocked in 5% BSA for 30 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (1/50) for 30 minutes at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen.

Tissues were counterstained with hematoxylin and mounted with DPX.




Fig3:; Immunohistochemical analysis of paraffin-embedded human liver carcinoma tissue using anti-NDUFB8 antibody. The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 8.0-8.4) for 20 minutes. The tissues were blocked in 5% BSA for 30 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (1/50) for 30 minutes at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




Fig4:; Immunohistochemical analysis of paraffin-embedded human kidney tissue using anti-NDUFB8 antibody. The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 8.0-8.4) for 20 minutes. The tissues were blocked in 5% BSA for 30 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (1/50) for 30 minutes at room temperature. The

detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




Fig5:; Immunohistochemical analysis of paraffin-embedded human small intestine tissue using anti-NDUFB8 antibody. The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 8.0-8.4) for 20 minutes. The tissues were blocked in 5% BSA for 30 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (1/50) for 30 minutes at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




Fig6:; Flow cytometric analysis of NDUFB8 was done on HeLa cells. The cells were fixed, permeabilized and stained with the primary antibody (1/50) (purple). After incubation of the primary antibody at room temperature for an hour, the cells were stained with a Alexa Fluor 488-conjugated Goat anti-Rabbit IgG Secondary

Anti-NDUFB8 antibody



Catalog Number: 176582

antibody at 1/1000 dilution for 30
minutes. Unlabelled sample was used as a control

(cells without incubation with primary antibody;
yellow).