

Anti-TNFRSF9 antibody



Catalog Number: 176678

General Information

Product name:

Anti-TNFRSF9 antibody

Immunogen:

Applications:

WB, IHC-P, ICC, FC

Specificity:

Human, Mouse, Rat

Antibody description:

Mouse monoclonal antibody to TNFRSF9

Properties

Preparation:

This antigen of this antibody was recombinant protein

Formulation:

Liquid, 1*PBS (pH7.4), 0.2% BSA, 50% Glycerol.
Preservative: 0.05% Sodium Azide.

Storage:

Store at +4°C after thawing. Aliquot store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.

Clonality:

Monoclonal

Ig Type: IgG1

Applications

WB: 1:500

ICC: 1:100-1:500

IHC-P: 1:50-1:200

FC: 1:50-1:200

Validations




Fig1: Western blot analysis of CD137 on HepG2 cell lysate using anti-CD137 antibody at 1/500 dilution.




Fig2: ICC staining CD137 (green) in A431 cells. The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.




Fig3: ICC staining CD137 (green) in A549 cells. The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.




Fig4: ICC staining CD137 (green) in LOVO cells. The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.




Fig5: Immunohistochemical analysis of paraffin-embedded rat brain tissue using anti-CD137 antibody. Counter stained with hematoxylin.




Fig6: Immunohistochemical analysis of paraffin-embedded rat hippocampus tissue using anti-CD137 antibody. Counter stained with hematoxylin.




Fig7: Immunohistochemical analysis of paraffin-embedded human tonsil tissue using anti-CD137 antibody. Counter stained with hematoxylin.





Fig8: Immunohistochemical analysis of paraffin-embedded human liver cancer tissue using anti-CD137 antibody. Counter stained with hematoxylin.



Catalog Number: 176678

Fig9: Immunohistochemical analysis of paraffin-embedded mouse brain tissue using anti-CD137 antibody. Counter stained with hematoxylin.




Fig10: Flow cytometric analysis of Jurkat cells with CD137 antibody at 1/100 dilution (red) compared with an unlabelled control (cells without incubation with primary antibody; black).