# Recombinant PODO protein (His tag)

| | |
|---|---|
| **Catalog Number** | 539017 |
| **Species** | Human |
| **Expression System** | E. coli |
| **Accession** | Q9NP85 |
| **Molecular Weight** | Predictes a molecular mass of 29.13 kDa. In SDS-PAGE under reducing conditions, it migrates as an approximately 29 kDa band. |

## Product Information

| | |
|---|---|
| **Sequence Region** | 100-342 |
| **Form** | Lyophilized powder |
| **Formulation** | Lyophilized from sterile PBS, pH 7.4 |
| **Purity** | >86%, by SDS-PAGE with Coomassie Brilliant Blue staining. |
| **Endotoxin** | Not tested. |
| **Activity** | Not tested. |
| **Stability** | Samples are stable for up to twelve months from date of receipt at -70℃ |
| **Protein Construction** | DNA encoding 100-342 aa (Q9NP85) were fused with 6His tag. |

## Storage & Handling

| | |
|---|---|
| **Reconstitution** | Reconstitute at 0.25 μg/μl in sterile water for short-term storage. Reconstitution with 50% glycerol solution is recommended for longer term storage (see Stability and Storage for more details). If a different concentration is needed for your purposes please adjust the reconstitution volume as required (please note: the ion concentration of the final solution will vary according to the volume used). Note: Centrifuge vial before opening. When reconstituting, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. |
| **Storage** | Store it under sterile conditions at -20℃ to -80℃. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles. |
| **Shipping** | In general, recombinant proteins are provided as lyophilized powder which are shipped at ambient temperature.<br>Bulk packages of recombinant proteins are provided as frozen liquid. They are shipped out with blue ice unless customers require otherwise. |

## Description

| | |
|---|---|
| **Description** | Recombinant Podocin (NPHS2) Fragment (aa 100-342) – a high-quality, stable protein fragment designed for structural and biochemical research. This 243-amino acid fragment offers researchers a soluble and functional unit ideal for studying glomerular permeability mechanisms.<br>Podocin, encoded by the NPHS2 gene, plays a critical role in regulating glomerular permeability by acting as a linker between the plasma membrane and cytoskeleton. In its full-length form, isoform 1 localizes to the cell membrane as a peripheral membrane protein, while isoform 2 is found in the endoplasmic reticulum. These localization patterns underscore Podocin's importance in maintaining renal filtration barrier integrity and its implications in podocyte function and kidney health. |