

Gene Name Synonym

PRG-1; 15-oxoprostaglandin 13-reductase; Dithiolethione-inducible gene 1 protein; D3T-inducible gene 1 protein; DIG-1; Leukotriene B4 12-hydroxydehydrogenase; NAD(P)H-dependent alkenal/one oxidoreductase

Protein Construction

DNA encoding 1-329 aa (Q14914) were fused with 6His tag.

Organism

Human

Expression Host

E. coli

Activity

Not tested.

Purity

>86%, by SDS-PAGE with Coomassie Brilliant Blue staining.

Endotoxin

Please contact the lab for more information.

Stability

Samples are stable for up to twelve months from date of receipt at -70°C

Molecular Mass

Predictes a molecular mass of 37.53 kDa. In SDS-PAGE under reducing conditions, it migrates as an approximately 38 kDa band.

Formulation

Lyophilized from sterile PBS, pH 7.4

Storage

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Reconstitute at 0.25 µg/µl in sterile water for short-term storage. Reconstitution with 50% glycerol solution is recommended for longer term storage (see Stability and Storage for more details). If a different concentration is needed for your purposes please adjust the reconstitution volume as required (please note: the ion concentration of the final solution will vary according to the volume used). Note: Centrifuge vial before opening. When reconstituting, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution.