

General Information

Gene Name Synonym

Cachectin; TNF-alpha; Tumor necrosis factor ligand superfamily member 2; Tumor necrosis factor, membrane form; N-terminal fragment; Intracellular domain 1; Intracellular domain 2; C-domain 1; C-domain 2; Tumor necrosis factor, soluble form

Protein Construction

A DNA sequence encoding the soluble form of mouse TNF- α (NP_038721.1) (Leu 80-Leu 235) was expressed, with an initial Met at the C-terminus.

Organism

Mouse

Expression Host

E. coli

QC Testing

Activity

Measured in a cytotoxicity assay using L929 mouse fibrosarcoma cells in the presence of the metabolic inhibitor actinomycin D. The ED_{50} for this effect is typically 0.04-0.08 ng/mL.

Purity

> 98 % as determined by SDS-PAGE

Endotoxin

Please contact us for more information.

Stability

Samples are stable for up to twelve months from date of receipt at -70°C

Predicted N terminal

Met

Molecular Mass

The recombinant mouse TNF- α consists of 157 amino acids and migrates with an apparent molecular mass of 17 kDa as estimated in SDS-PAGE under reducing conditions.

Formulation

Lyophilized from sterile 50mM Tris 0.2M NaCl, pH 7.2

1. 5 % trehalose and mannitol are added as protectants before lyophilization.

2. Please contact us for any concerns or special requirements.

Usage Guide


Storage

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/mL. Concentration is measured by UV-Vis.

SDS-PAGE



Mouse TNF-alpha / TNFA Protein SDS-PAGE