Catalog Number: 500661



General Information

Gene Name Synonym

Neuregulin-1; Acetylcholine receptor-inducing activity; Breast cancer cell differentiation factor p45; Glial growth factor; Heregulin; Neu differentiation factor; Sensory and motor neuronderived factor

Protein Construction

A DNA sequence encoding the N-terminal fragment (Ser 2-Lys 246) of human NRG1 isoform beta1 (Q02297-6) was expressed.

Organism

Human

Expression Host

Human Cells

QC Testing

Activity

Measured in a cell proliferation assay using MCF-7 cells. The ED_{50} for this effect is typically 2-10 ng/mL.

Purity

> 90 % as determined by SDS-PAGE

Endotoxin

< 1.0 EU per μg of the protein as determined by the LAL method

Stability

Samples are stable for up to twelve months from date of receipt at -70 $^\circ \rm C$

Predicted N terminal

Ser 2 Molecular Mass

The recombinant human NRG1 comprises 245 amino acids and has a predicted molecular mass of 26.8 kDa.

Formulation

Lyophilized from sterile PBS, pH 7.41. 5 % trehalose and mannitol are added as protectants before lyophilization.2. Please contact us for any concerns or special requirements.

Usage Guide

Storage

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE

KDa	M
116	
66.2	-
45.0	
35.0	-
25.0	-
18.4	-
14.4	_

Human NRG1-beta 1 Protein () SDS-PAGE