Catalog Number: 502027



General Information

Gene Name Synonym

Neural apoptosis-regulated convertase 1; Proprotein convertase 9; Subtilisin/kexin-like protease PC9

Protein Construction

A DNA sequence encoding the human proprotein convertase 9 (AAV67948.1) (Met 1-Gln 692) was expressed with a C-terminal polyhistidine tag.

Organism

Human

Expression Host

Human Cells

QC Testing

Activity

Measure by its ability to bind with human LDLR in a functional ELISA. Immobilized recombinant human PCSK9 at 10 μ g/ml (100 μ l/well) can bind biotinylated human LDLR. The EC₅₀ of biotinylated human LDLR is 0.61 μ g/ml.

Purity

> 97 % as determined by SDS-PAGE

Endotoxin

< 1.0 EU per μg of the protein as determined by the LAL method

Stability

Samples are stable for up to twelve months from date of receipt at -70 $^{\circ}\mathrm{C}$

Predicted N terminal

Gln 31

Molecular Mass

The recombinant human PCSK9 consists of 662 amino acids and predicts a molecular mass of 72.6 kDa. As a result of glycosylation and proteolytic digestion, rhPCSK9 migrates as doublet with apparent molecular mass of 20 kDa and 62 kDa respectively in SDS-PAGE under reducing conditions.

Formulation

Lyophilized from sterile PBS, pH 7.41. 5 % trehalose and mannitol are added as protectants before lyophilization.2. Please contact us for any concerns or special requirements.

Usage Guide

Storage

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE

KDa	М
116	
66.2	-
45.0	-
35.0	-
25.0	-
18.4	
14.4	-

Human PCSK9 / NARC1 Protein (His Tag) SDS-PAGE