Catalog Number: 502099



General Information

Gene Name Synonym

Activin-binding protein

Protein Construction

A DNA sequence encoding the mouse FST (P47931) (Met 1-Asn 317) was expressed with the Fc region of human IgG1 at the C-terminus.

Organism

Mouse

Expression Host

CHO Stable Cells

QC Testing

Activity

Measured by its ability to neutralize Activinmediated inhibition on MPC11 cell proliferation. The ED_{50} for this effect is typically 40-200 ng/mL in the presence of 10 ng/mL rhActivin A.

Purity

> 90 % as determined by SDS-PAGE

Endotoxin

< 1.0 EU per μg of the protein as determined by the LAL method

Stability

Samples are stable for up to twelve months from date of receipt at -70 $^{\circ}\mathrm{C}$

Predicted N terminal

Gly 30

Molecular Mass

The recombinant mouse FST comprises 528 amino acids and has a predicted molecular mass of 58.4 kDa. In SDS-PAGE under reducing conditions, the apparent molecular mass of the protein is approximately 66 kDa due to glycosylation.

Formulation

Lyophilized from sterile PBS, pH 7.4 1.5 % trehalose and mannitol are added as protectants before lyophilization. 2. Please contact us for any concerns or special requirements.

Usage Guide

Storage

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE

KDa M 116	-
66.2	-
45.0	
35.0	-
25.0	
18.4	1
14.4	

Mouse Follistatin / FST (FS288) Protein (Fc Tag) SDS-PAGE