

General Information

Gene Name Synonym

Protein Cd40; Tumor necrosis factor receptor superfamily, member 5, isoform CRA_a

Protein Construction

A DNA sequence encoding the rat CD40 (Q4QQW2) (Met1-Arg193) was expressed, fused with the Fc region of human IgG1 at the C-terminus.

Organism

Rat

Expression Host

Human Cells

QC Testing

Purity

> 95 % as determined by SDS-PAGE

Endotoxin

< 1.0 EU per μ g of the protein as determined by the LAL method

Stability

Samples are stable for up to twelve months from date of receipt at -70°C

Predicted N terminal

Leu 20

Molecular Mass

The recombinant rat CD40/Fc is a disulfide-linked homodimer. The reduced monomer comprises 415 amino acids and has a predicted molecular mass of 46.2 kDa. The apparent molecular mass of the protein is approximately 53 kDa in SDS-PAGE under reducing conditions.

Formulation

Lyophilized from sterile PBS, pH 7.4

1. 5 % trehalose and mannitol are added as protectants before lyophilization.

2. Please contact us for any concerns or special requirements.

Usage Guide


Storage

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE



Rat CD40 / TNFRSF5 Protein (Fc Tag) SDS-PAGE