

General Information

Gene Name Synonym

Osteoblast-specific factor 2

Protein Construction

A DNA sequence encoding the human POSTN isoform 3 (NP_001129407.1) corresponding to amino acid (Met 1-Gln 781) was expressed with a C-terminal polyhistidine tag.

Organism

Human

Expression Host

Human Cells

QC Testing

Activity

Measured by its ability to induce adhesion of ATDC5 mouse chondrogenic cells. When cells are added to POSTN-His coated plates (10µg/mL, 100µL/well), approximately >60% will adhere specifically after 30 minutes at 37°C.

Purity

> 97 % as determined by SDS-PAGE

Endotoxin

< 1.0 EU per µg of the protein as determined by the LAL method

Stability

Samples are stable for up to twelve months from date of receipt at -70°C

Predicted N terminal

Asn 22

Molecular Mass

The recombinant human POSTN consists of 771 amino acids and has a predicted molecular mass of 86 kDa. In SDS-PAGE under reducing conditions, it migrates with an apparent molecular mass of 80 kDa due to glycosylation.

Formulation

Lyophilized from sterile PBS, pH 7.4

1. 5 % trehalose and mannitol are added as protectants before lyophilization.

2. Please contact us for any concerns or special requirements.

Usage Guide


Storage

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE



Human Periostin / POSTN Protein (His Tag) SDS-PAGE