

Catalog Number: 502785

General Information

Gene Name Synonym

C-X-C motif chemokine 7; Leukocyte-derived growth factor; Macrophage-derived growth factor; Small-inducible cytokine B7; Connective tissueactivating peptide III; LA-PF4; Low-affinity platelet factor IV; TC-2; Connective tissueactivating peptide III(1-81); Beta-thromboglobulin; Neutrophil-activating peptide 2(74); Neutrophilactivating peptide 2(73); Neutrophil-activating peptide 2; TC-1; Neutrophil-activating peptide 2(1-66); Neutrophil-activating peptide 2(1-63)

Protein Construction

A DNA sequence encoding the human TPK1 (P02775) (Met1-Ser243) was expressed with a polyhistidine tag at the N-terminus.

Organism

Human

Expression Host

E. coli

QC Testing

Activity

Kinase activity untested

Purity

> 90 % as determined by SDS-PAGE

Endotoxin

Please contact us for more information.

Stability

Samples are stable for up to twelve months from date of receipt at -70 $^{\circ}\mathrm{C}$

Predicted N terminal His

Molecular Mass

The recombinant human TPK1 consists of 261 amino acids and predicts a molecular mass of 29.5 KDa. It migrates as an approximately 27-32 KDa band in SDS-PAGE under reducing conditions.

Formulation

Lyophilized from sterile PBS, pH 7.4.1. 5 % trehalose and mannitol are added as protectants before lyophilization.2. Please contact us for any concerns or special requirements.

Usage Guide

Storage

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE

KDa	M
116	
66.2	-
45.0	-
35.0	-
25.0	-
18.4	-
14.4	-

Human TPK1 Protein (His Tag) SDS-PAGE