

Human IL12A / IL-12A (Fc Tag) recombinant protein



Catalog Number: 503245

General Information

Gene Name Synonym:

Cytotoxic lymphocyte maturation factor 35 kDa subunit; IL-12 subunit p35; NK cell stimulatory factor chain 1

Protein Construction:

A DNA sequence encoding the p35 subunit of human IL12, termed as IL12A (P29459) (Met 1-Ser 219) was fused with the Fc region of human IgG1 at the C-terminus.

Source: Human

Expression Host: Human Cells

QC Testing

Activity:

Purity:

> 95 % as determined by SDS-PAGE

Endotoxin:

< 1.0 EU per μ g of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70°C

Predicted N terminal:

Arg 23

Molecular Mass:

The recombinant human IL12A/Fc is a disulfide-linked homodimer. The reduced monomer consists

of 435 amino acids and has a predicted molecular mass of 49.3 kDa. As a result of glycosylation, the rhIL12A/Fc monomer migrates as an approximately 65-70 kDa band in SDS-PAGE under reducing conditions.

Formulation:

Lyophilized from sterile PBS, pH 7.4

1. 5 % trehalose and mannitol are added as protectants before lyophilization.

2. Please contact us for any concerns or special requirements.

Usage Guide


Storage:

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution:

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE:



Human IL12A / NKSF1 Protein (Fc Tag) SDS-PAGE