

Human NAP-2 / PPBP / CXCL7 (His Tag) recombinant protein



Catalog Number: 503598

General Information

Gene Name Synonym

C-X-C motif chemokine 7; Leukocyte-derived growth factor; Macrophage-derived growth factor; Small-inducible cytokine B7; Connective tissue-activating peptide III; LA-PF4; Low-affinity platelet factor IV; TC-2; Connective tissue-activating peptide III(1-81); Beta-thromboglobulin; Neutrophil-activating peptide 2(74); Neutrophil-activating peptide 2(73); Neutrophil-activating peptide 2; TC-1; Neutrophil-activating peptide 2(1-66); Neutrophil-activating peptide 2(1-63)

Protein Construction

A DNA sequence encoding the human CXCL7 (P02775) (Ala59-Asp128) was expressed with a polyhistidine tag at the N-terminus.

Organism

Human

Expression Host

E. coli

QC Testing

Purity

> 90 % as determined by SDS-PAGE

Endotoxin

Please contact us for more information.

Stability

Samples are stable for up to twelve months from date of receipt at -70°C

Predicted N terminal

His

Molecular Mass

The recombinant human CXCL7 consists of 88 amino acids and predicts a molecular mass of 9.9 KDa. It migrates as an approximately 13 KDa band in SDS-PAGE under reducing conditions.

Formulation

Lyophilized from sterile PBS, pH 7.4.

1. 5 % trehalose and mannitol are added as protectants before lyophilization.

2. Please contact us for any concerns or special requirements.

Usage Guide


Storage

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE



Human NAP-2 / PPBP / CXCL7 Protein (His Tag)
SDS-PAGE