Human CD32a / Fc gamma RIIA / FCGR2A (167 Arg, His Tag) recombinant protein

Catalog Number: 504175

General Information

Gene Name Synonym

CDw32; Fc-gamma RII-a

Protein Construction

A DNA sequence encoding (Met 1-Ile 218) of human CD32a (AAA35827.1) was expressed with a C-terminal polyhistidine tag. It is identical to FCGR2A131H/R in the reference.

Organism

Human

Expression Host

CHO Cells

QC Testing

Activity

Measured by its binding ability in a functional ELISA.

Immobilized human CD32a-His (Cat:504175) at 10 μ g/ml (100 μ l/well) can bind biotinylated human IgG1, The EC₅₀ of biotinylated human IgG1 is 0.14-0.32 μ g/ml.

Purity

> 97 % as determined by SDS-PAGE

Endotoxin

< 1.0 EU per μg of the protein as determined by the LAL method

Stability

Samples are stable for up to twelve months from date of receipt at -70 $^{\circ}\mathrm{C}$

Predicted N terminal

Gln 34 **Molecular Mass**

The secreted recombinant human CD32a consists of 196 amino acids and predicts a molecular mass of 22 KDa. The apparent molecular mass of the protein is approximately 29-32 KDa in SDS-PAGE under reducing conditions due to glycosylation.

Formulation

Lyophilized from sterile PBS, pH 7.4 1.5 % trehalose and mannitol are added as protectants before lyophilization. 2. Please contact us for any concerns or special requirements.

Usage Guide

Storage

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE

| KDa | м |
|------|---|
| 116 | |
| 66.2 | |
| 45.0 | |
| 35.0 | - |
| 25.0 | _ |
| 18.4 | - |
| 14.4 | |

Human CD32a / FCGR2A Protein (167 Arg, His Tag) SDS-PAGE

