

Human FGFR2/CD332 (His & Fc Tag) recombinant protein



Catalog Number: 504217

General Information

Gene Name Synonym

K-sam; Keratinocyte growth factor receptor

Protein Construction

A DNA sequence encoding the human FGFR2 (NP_000132.3) extracellular domain (Met 1-Glu 377) was fused with the C-terminal polyhistidine-tagged Fc region of human IgG1 at the C-terminus.

Organism

Human

Expression Host

Human Cells

QC Testing

Activity

Measured by its ability to inhibit FGF-acidic (aFGF/FGF1) dependent proliferation of Balb/C 3T3 mouse fibroblasts.
The ED₅₀ for this effect is typically 0.5-2.5 ng/ml.

Purity

> 90 % as determined by SDS-PAGE

Endotoxin

< 1.0 EU per µg of the protein as determined by the LAL method

Stability

Samples are stable for up to twelve months from date of receipt at -70°C

Predicted N terminal

Arg 22

Molecular Mass

The recombinant human FGFR2/Fc is a disulfide-linked homodimer after removal of the signal peptide. The reduced monomer consists of 604 amino acids and has a predicted molecular mass of 67.6 kDa. In SDS-PAGE under reducing conditions, the apparent molecular mass of rhFGFR2/Fc monomer is approximately 110-120 kDa due to glycosylation.

Formulation

Lyophilized from sterile PBS, pH 7.4

1. 5 % trehalose and mannitol are added as protectants before lyophilization.
2. Please contact us for any concerns or special requirements.

Usage Guide


Storage

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE



Human FGFR2 Protein (His & Fc Tag) SDS-PAGE