

# Human 2B4 / CD244 (His Tag) recombinant protein



Catalog Number: 504221

## General Information

### Gene Name Synonym

NK cell activation-inducing ligand; NK cell type I receptor protein 2B4; SLAM family member 4; Signaling lymphocytic activation molecule 4

### Protein Construction

A DNA sequence encoding the extracellular domain (Met 1-Arg 221) of human 2B4 (NP\_057466.1) was expressed, with a polyhistidine tag at the C-terminus.

### Organism

Human

### Expression Host

Human Cells

## QC Testing

### Activity

Measured by its binding ability in a functional ELISA. Immobilized human 2B4 at 2 µg/ml (100 µl/well) can bind human CD48. The  $EC_{50}$  of human CD48 is 0.39 µg/ml.

### Purity

> 95 % as determined by SDS-PAGE

### Endotoxin

< 1.0 EU per µg of the protein as determined by the LAL method

### Stability

Samples are stable for up to twelve months from date of receipt at -70°C

### Predicted N terminal

Cys 22

### Molecular Mass

The recombinant 2B4 comprises 211 amino acids and predicts a molecular mass of 23.8 kDa. As a result of glycosylation, the rh 2B4 protein migrates as an approximately 45-50 kDa band in SDS-PAGE under reducing conditions.

### Formulation

Lyophilized from sterile PBS, pH 7.4

1. 5 % trehalose and mannitol are added as protectants before lyophilization.

2. Please contact us for any concerns or special requirements.

## Usage Guide


### Storage

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

### Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

### SDS-PAGE



Human 2B4 / SLAMF / CD244 Protein (His Tag)  
SDS-PAGE