

General Information

Protein Construction

A DNA sequence encoding the rhesus CD1C(NP_001139005.1) (Met1-Phe298) of was expressed with a polyhistidine tag at the C-terminus, constructed the plasmid 1; A DNA sequence encoding the rhesus B2M (NP_001040602.1) (Met1-Met119) constructed the plasmid 2. The two plasmids were co-expressed and the CD1C/B2M heterodimer was purified.

Organism

Rhesus

Expression Host

Human Cells

QC Testing

Purity

> (64.1+32.5) % as determined by SDS-PAGE.

Endotoxin

<1.0 EU per μg protein as determined by the LAL method.

Stability

Samples are stable for up to twelve months from date of receipt at -70°C

Predicted N terminal

Gln21 & Ile21

Molecular Mass

The recombinant heterodimer of rhesus CD1C&B2M comprises 388 (289+99) amino acids and has a calculated molecular mass of 44.7 (33.1+ 11.6) KDa.

Formulation

Lyophilized from sterile PBS, pH 7.4.

1. 5 % trehalose and mannitol are added as protectants before lyophilization.
2. Please contact us for any concerns or special requirements.

Usage Guide


Storage

Store it under sterile conditions at -20°C to -80°C . It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE



Rhesus CD1C & B2M Heterodimer Protein