pMAl-c2E

Vector ID	V006980
Antibiotic Resistance	Ampicillin
Length	6648 bp
Vector Type	E. coli Expression Vectors

Product Information

Replication Origin	ori
Copy Number	High copy number
Promoter	Tac
Cloning Method	Enzyme digestion and ligation
5' Primer	MalE: 5'-GGTCGTCAGACTGTCGATGAAGCC-3'; MBP-F: 5'-gatgaagccctgaaagacgcgcag-3'
3' Primer	pBad-R: 5'-gatttaatctgtatcagg-3'; M13-F: 5'-TGTAAAACGACGGCCAGT-3'
Fusion Tag	N-MBP, N-Factor Xa
Growth Strain(s)	Top10
Comment	pMAl-c2E is a plasmid expressing proteins in the cytoplasm as fusions to maltose-binding protein.

Vector Map



Plasmid Resuspension Protocol

1. Centrifuge at 5,000×g for 5 min.
2. Carefully open the tube and add sterile water to dissolve the DNA: add 20 µl for 5 µg plasmid, and 100 µl for 100 µg plasmid.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin to concentrate the liquid at the bottom. Speed is less than 5000×g.
5. Store the plasmid at -20 ℃.