Anti-CD38 antibody

Catalog Number: 102060



Product name

Anti-CD38 antibody

Specificity

Mouse CD38

Antibody description

Rabbit polyclonal to CD38

Preparation

Produced in rabbits immunized with purified, recombinant Mouse CD38 (rM CD38; NP_031672.2; Leu 45-Thr 304). CD38 specific IgG was purified by mouse CD38 affinity chromatography

Formulation

0.2 µm filtered solution in PBS with 5% trehalose

Storage

This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C.

Preservative-Free.

Sodium azide is recommended to avoid contamination (final concentration 0.05%-0.1%). It is toxic to cells and should be disposed of properly. Avoid repeated freeze-thaw cycles.

Clonality

Polyclonal

Ig Type

Rabbit IgG

Applications

ELISA, IHC-P, FCM

Dilutions


ELISA: 0.5-1.0 μg/mL

This antibody can be used at 0.5-1.0 μ g/mL with the appropriate secondary reagents to detect Mouse CD38. The detection limit for Mouse CD38 is 0.00245 ng/well.

IHC-P: $0.1-2 \mu g/mL$


FCM: 0.5-2 µg/Test

Validations



CD38 Antibody, Rabbit PAb, Antigen Affinity Purified, Immunohistochemistry

Immunochemical staining of mouse CD38 in mouse kidney with rabbit polyclonal antibody (0.2 μ g/mL, formalin-fixed paraffin embedded sections). Positive staining was localized to



endotheliocyte.

CD38 Antibody, Rabbit PAb, Antigen Affinity Purified, Flow cytometric

Flow cytometric analysis of CD38 expression on spleen lymphocytes. BALB/c splenocytes were


Anti-CD38 antibody

Catalog Number: 102060



stained with Mouse CD38 Affinity Purified Polyclonal Ab (50191-RP02, 1 µg/test), then a FITC-conjugated second step antibody. The histogram were derived from the gated events based on light scattering characteristics of

lymphocytes.



CD38 Antibody, Rabbit PAb, Antigen Affinity Purified, Immunohistochemistry

Immunochemical staining of mouse CD38 in mouse heart with rabbit polyclonal antibody (0.2 $\,$ µg/mL, formalin-fixed paraffin embedded sections). Positive staining was localized to endotheliocyte.