# Recombinant HYCC2 protein (His tag)

| | |
|---|---|
| **Catalog Number** | 549571 |
| **Species** | Human |
| **Expression System** | E. coli |
| **Accession** | Q8IXS8 |
| **Molecular Weight** | Predictes a molecular mass of 35.06 kDa. In SDS-PAGE under reducing conditions, it migrates as an approximately 35 kDa band. |

## Product Information

| | |
|---|---|
| **Sequence Region** | 1-291 |
| **Form** | Lyophilized powder |
| **Formulation** | Lyophilized from sterile PBS, pH 7.4 |
| **Purity** | >87%, by SDS-PAGE with Coomassie Brilliant Blue staining. |
| **Endotoxin** | Not tested. |
| **Activity** | Not tested. |
| **Stability** | Samples are stable for up to twelve months from date of receipt at -70℃ |
| **Protein Construction** | DNA encoding 1-291 aa (Q8IXS8) were fused with 6His tag. |

## Storage & Handling

| | |
|---|---|
| **Reconstitution** | Reconstitute at 0.25 μg/μl in sterile water for short-term storage. Reconstitution with 50% glycerol solution is recommended for longer term storage (see Stability and Storage for more details). If a different concentration is needed for your purposes please adjust the reconstitution volume as required (please note: the ion concentration of the final solution will vary according to the volume used). Note: Centrifuge vial before opening. When reconstituting, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. |
| **Storage** | Store it under sterile conditions at -20℃ to -80℃. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles. |
| **Shipping** | In general, recombinant proteins are provided as lyophilized powder which are shipped at ambient temperature.<br>Bulk packages of recombinant proteins are provided as frozen liquid. They are shipped out with blue ice unless customers require otherwise. |

## Description

| | |
|---|---|
| **Description** | Recombinant Hyccin 2 (HYCC2) fragment (amino acids 1-291) from Homo sapiens, designed for structural and biochemical studies. This fragment represents a stable, soluble unit of the protein, facilitating investigations into its role in phosphatidylinositol 4-kinase (PI4K) localization at the plasma membrane. Hyccin 2, encoded by the HYCC2 gene in humans, functions as a component of a complex essential for localizing phosphatidylinositol 4-kinase (PI4K) to the plasma membrane, a critical process in cellular signaling and membrane dynamics. The protein is primarily located in the cytoplasm and cytosol, with additional presence at the cell membrane, underscoring its involvement in membrane-associated regulatory mechanisms. This recombinant fragment (1-291) enables focused research on these functional attributes without domain-specific constraints. |