# **Anti-THEMIS antibody**

Catalog Number: 175723



#### Product name

Anti-THEMIS antibody

# **Specificity**

Human, Mouse, Rat, Dog, Pig, Horse, Rabbit

### **Antibody description**

Rabbit polyclonal antibody to THEMIS

## **Preparation**

This antigen of this antibody was klh conjugated synthetic peptide derived from human themis 301-400/641

#### **Formulation**

Liquid, 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

#### **Storage**

Store at -20°C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4°C.

#### **Clonality**

Polyclonal

#### Ig Type

Rabbit IgG

#### **Applications**

WB, IHC-P

#### **Dilutions**

WB:1:500-2000

IHC-P:1:400-800

#### **Validations**





Fig1: Tissue/cell: human kidney tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min;; Incubation: Anti-THEMIS Polyclonal Antibody, Unconjugated 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010)



staining

Fig2: Tissue/cell: human kidney tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;; Antigen retrieval: citrate buffer ( 0.01M, pH 6.0 ), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;; Incubation: Anti-THEMIS Polyclonal Antibody, Unconjugated 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010)

# **Anti-THEMIS antibody**

Catalog Number: 175723






Fig3: Sample:; U937 Cell (Human) Lysate at 30 ug; Primary: Anti-THEMIS at 1/300 dilution; Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution; Predicted band size: 73 kD;

Observed band size: 73 kD